

# POLIO VACCINE

## WHAT YOU NEED TO KNOW

### 1 What is polio?

**Polio is a disease caused by a virus.** It enters a child's (or adult's) body through the mouth. Sometimes it does not cause serious illness. But sometimes it causes *paralysis* (can't move arm or leg). It can kill people who get it, usually by paralyzing the muscles that help them breathe.

Polio used to be very common in the United States. It paralyzed and killed thousands of people a year before we had a vaccine for it.

### 2 Why get vaccinated?

**Inactivated Polio Vaccine (IPV) can prevent polio.**

**History:** A 1916 polio epidemic in the United States killed 6,000 people and paralyzed 27,000 more. In the early 1950's there were more than 20,000 cases of polio each year. **Polio vaccination was begun in 1955.** By 1960 the number of cases had dropped to about 3,000, and by 1979 there were only about 10. The success of polio vaccination in the U.S. and other countries sparked a world-wide effort to eliminate polio.

**Today:** No wild polio has been reported in the United States for over 20 years. But the disease is still common in some parts of the world. It would only take one case of polio from another country to bring the disease back if we were not protected by vaccine. If the effort to eliminate the disease from the world is successful, some day we won't need polio vaccine. Until then, we need to keep getting our children vaccinated.

#### **Oral Polio Vaccine: No longer recommended**

There are two kinds of polio vaccine: **IPV**, which is the shot recommended in the United States today, and a live, oral polio vaccine (**OPV**), which is drops that are swallowed.

Until recently OPV was recommended for most children in the United States. OPV helped us rid the country of polio, and it is still used in many parts of the world.

Both vaccines give immunity to polio, but OPV is better at keeping the disease from spreading to other people. However, for a few people (about one in 2.4 million), OPV actually causes polio. Since the risk of getting polio in the United States is now extremely low, experts believe that using oral polio vaccine is no longer worth the slight risk, except in limited circumstances which your doctor can describe. The polio shot (IPV) does not cause polio. **If you or your child will be getting OPV, ask for a copy of the OPV supplemental Vaccine Information Statement.**

### 3 Who should get polio vaccine and when?

IPV is a shot, given in the leg or arm, depending on age. Polio vaccine may be given at the same time as other vaccines.

#### **Children**

Most people should get polio vaccine when they are children. Children get 4 doses of IPV, at these ages:

- ✓ A dose at 2 months
- ✓ A dose at 4 months
- ✓ A dose at 6-18 months
- ✓ A booster dose at 4-6 years

#### **Adults**

Most adults do not need polio vaccine because they were already vaccinated as children. But three groups of adults are at higher risk and *should* consider polio vaccination:

- (1) people traveling to areas of the world where polio is common,
- (2) laboratory workers who might handle polio virus, and
- (3) health care workers treating patients who could have polio.

Adults in these three groups who **have never been vaccinated against polio** should get 3 doses of IPV:

- ✓ The first dose at any time,
- ✓ The second dose 1 to 2 months later,
- ✓ The third dose 6 to 12 months after the second.

Adults in these three groups who **have had 1 or 2 doses** of polio vaccine in the past should get the remaining 1 or 2 doses. It doesn't matter how long it has been since the earlier dose(s).

Adults in these three groups who **have had 3 or more doses** of polio vaccine (either IPV or OPV) in the past may get a booster dose of IPV.

Ask your health care provider for more information.

## 4

### Some people should not get IPV or should wait.

#### These people should not get IPV:

- Anyone who has ever had a life-threatening allergic reaction to the antibiotics **neomycin**, **streptomycin** or **polymyxin B** should not get the polio shot.
- Anyone who has a severe allergic reaction to a polio shot should not get another one.

#### These people should wait:

- Anyone who is moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting polio vaccine. People with minor illnesses, such as a cold, *may* be vaccinated.

Ask your health care provider for more information.

## 5

### What are the risks from IPV?

Some people who get IPV get a sore spot where the shot was given. The vaccine used today has never been known to cause any serious problems, and most people don't have any problems at all with it.

However, a vaccine, like any medicine, could cause serious problems, such as a severe allergic reaction. *The risk of a polio shot causing serious harm, or death, is extremely small.*

## 6

### What if there is a serious reaction?

#### What should I look for?

Look for any unusual condition, such as a serious allergic reaction, high fever, or unusual behavior.

If a serious allergic reaction occurred, it would happen within a few minutes to a few hours after the shot. Signs of a serious allergic reaction can include difficulty breathing, weakness, hoarseness or wheezing, a fast heart beat, hives, dizziness, paleness, or swelling of the throat

#### What should I do?

- Call a doctor, or get the person to a doctor right away.

- Tell your doctor what happened, the date and time it happened, and when the vaccination was given.
- Ask your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS website at [www.vaers.org](http://www.vaers.org), or by calling 1-800-822-7967.

*VAERS does not provide medical advice.*

Reporting reactions helps experts learn about possible problems with vaccines.

## 7

### The National Vaccine Injury Compensation Program

In the rare event that you or your child has a serious reaction to a vaccine, there is a federal program that can help pay for the care of those who have been harmed.

For details about the National Vaccine Injury Compensation Program, call **1-800-338-2382** or visit the program's website at <http://www.hrsa.gov/osp/vicp>

## 8

### How can I learn more?

- Ask your doctor or nurse. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department's immunization program.
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)**
  - Visit the National Immunization Program's website at <http://www.cdc.gov/nip>



**U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES**  
Centers for Disease Control and Prevention  
National Immunization Program



# IMMUNIZE L.A.

## Polio Vaccination Consent Form

I have read and understood the above information about Polio vaccination. I have had a chance to ask questions. I understand the benefits and risks of Polio vaccination and request that vaccine be given to me or the person named below whom I am authorized to sign for.

Information of Person to Receive Vaccine			
Name (Please print)	Birth date	Age	
Address: Street	City	State	Zip
Phone Number			
X			
Signature (Person receiving Vaccine or Parent or Guardian)			Date

For Clinic Use	
<b>Plaza West Pharmacy</b> 19631 Parthenia Street Northridge, CA 91324 Phone: (818) 886-4900	<b>Olympia Plaza Pharmacy</b> 5901 W. Olympic Blvd., #103 Los Angeles, CA 90036 Phone: (323) 937-2590
<b>Date of Vaccination:</b>	
<b>Manufacture/ lot number:</b>	
<b>Site of Injection:</b>	
<b>Chronic Diseases:</b>	Yes                      No



# IMMUNIZE L.A.

## Screening Questionnaire

The following questions will help us determine which vaccines may be given today. If a question is not clear, please ask your health care provider to explain it.

- |   | Yes | No | Don't<br>Know |
|---|-----|----|---------------|
| 1. Are you sick today?  |     |    |               |
| 2. Do you have allergies to medications, food, or any vaccine?  |     |    |               |
| 3. Have you ever had a serious reaction after receiving a vaccination?  |     |    |               |
| 4. Do you have cancer, leukemia, AIDS, or any other immune system problem?  |     |    |               |
| 5. Do you take cortisone, prednisone, other steroids, or anticancer drugs, or have you had x-ray treatments?                                  |     |    |               |
| 6. During the past year, have you received a transfusion of blood or blood products, or been given a medicine called immune (gamma) globulin? |     |    |               |
| 7. For women: Are you pregnant or is there a chance you could become pregnant in the next three months?                                       |     |    |               |
| 8. Have you received any vaccinations in the past 4 weeks?  |     |    |               |

X

Signature (Person receiving Vaccine or Parent or Guardian)

Date

**Did you bring your immunization record card with you?**      Yes       No

It is important for you to have a personal record of your vaccinations. If you don't have a record card, ask your health care provider to give you one. Bring this record with you every time you seek medical care. Make sure your health care provider records all your vaccinations on it.

<b>For Clinic Use</b>	
<b>Plaza West Pharmacy</b> 19631 Parthenia Street Northridge, CA 91324 Phone: (818) 886-4900	<b>Olympia Plaza Pharmacy</b> 5901 W. Olympic Blvd., #103 Los Angeles, CA 90036 Phone: (323) 937-2590